

## PLUS TRAP · Attractants



### Description

**Plus Trap®** is a ready-to-use bait trap with a powerful attractant for *Ceratitis capitata* and *Anastrepha* spp which, through the emission of volatile compounds, exerts a powerful attraction, especially in females flies of the species, which are attracted to the interior of the trap and, as they are unable to escape, they die by drowning inside.

### Features

- ▶ **Specific.** **Plus Trap®** is a specific food attractant for *Ceratitis capitata* and *Anastrepha* spp with a high protein content that exerts a high power of attraction, especially in females flies.
- ▶ **Clean.** **Plus Trap®** is a biological product free of insecticides. Suitable for use in organic production.
- ▶ **Ready to use.** **Plus Trap®** it is not necessary to dilute or add other products.
- ▶ **Affordable.** **Plus Trap®** allows the re-baiting of the traps through its economical refill drum.
- ▶ **Contains trimethylamine** and protein **hydrolysate**.
- ▶ **Lifespan:** More than 30 days (depending on weather conditions)
- ▶ **Toxicity:** **Low**

### Packaging

Box containing 24 traps (960 traps per pallet of 1m x 1.20m x 2m).  
Economical 20-litre refill drum (metering valves are also available).

### Forma de Uso

Plus Trap® is ready to use. It is not necessary to add additives or dilute the product. You should remove the strip and hung the product from the tree. That's it.

The amount of Plus Trap® per hectare will depend on the pressure, dynamics of the pest and type of crop to be protected, with a minimum of 70 traps per hectare.

When setting up the traps, special attention should be paid to the dynamics of the pest within the property, reinforcing them in places where the greatest pressure is detected.

Hang Plus Trap® 1.5 meters above the ground in the sunny part of the plant; the holes of the trap should not be obstructed by leaves or branches.

Plus Trap® will need to be replenished before it fully evaporates. The time interval for its replacement may vary depending on temperature, humidity, exposure to sunlight, etc.

| Crops                                                                                                                                                                                                                                                                              | Pests                                                    | Doses                                                                                  | Application time                                                                                                                                                                                                                                      |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------|----------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Citric:</b> (Orange tree, mandarin orange tree, grapefruit tree)<br><br><b>Stone fruit trees:</b> (peach tree, plum tree, olive tree, apricot tree)<br><br><b>Seed fruit trees:</b> (pear tree, apple tree, vines)<br><br><b>Tropical fruit trees:</b> (mango tree, guava tree) | <b>Fruit flies:</b> (Ceratitis capitata, Anastrepha spp) | A minimum of 70 traps per hectare, depending on the pressure and dynamics of the pest. | The time, the concentration and the number of applications will be determined according to the thresholds of economic damage through attractive traps for the detection of adult flies. It is suggested to monitor these traps at least twice a week. |

## Preservation

Store in a cool and dry place, avoiding exposure to sunlight.

## Safety Regulations

Keep out of reach of children and inexperienced people. Handle with gloves. Wash the exposed surface with plenty of soap and water. If ingested, you should induce vomiting and seek medical advice immediately.

## Warnings

Avoid product spillages in order to prevent attraction areas outside the trap. Confer with an agricultural science professional before purchasing and using this product.

**Susbin S.A.** is responsible for the composition of the product until its expiration date.

**Susbin S.A.** is not responsible for damages of any nature that may derive from a different use of the product.