



Description

SWD Lure® is a bait trap ready to be used with a powerful attractant for *Drosophila suzukii* (especially females flies) which, through the emission of volatile compounds, exerts a powerful attraction on this pest, not requiring insecticides since they cannot leave the trap and die by drowning inside.

Features

- ▶ **Specific.** **SWD Lure®** is a specific food attractant for *Drosophila suzukii* that exerts a high power of attraction, especially in females flies.
- ▶ **Clean.** **SWD Lure®** is a biological product free of insecticides. Suitable for use in organic production.
- ▶ **Ready to use.** **SWD Lure®** it is not necessary to dilute or add other products.
- ▶ **Affordable.** **SWD Lure®** allows the re-baiting of the traps through its economical refill drum.
- ▶ **Peptides** 2,14% w/w
- ▶ **Inert organic acids** 86,99% w/w
- ▶ **q.s.:** 100% w/w
- ▶ **Lifespan:** More than 30 days (depending on weather conditions).
- ▶ **Toxicity:** **Low**

Packaging

- Economical 20-litre refill drum (metering valves are also available).
- Box containing 24 traps (960 traps per pallet of 1m x 1.20m x 2m).

Usage

SWD Lure® is ready to use. It is not necessary to add additives or dilute the product.

The amount of SWD Lure® per hectare will depend on the pressure, dynamics of the pest and type of crop to be protected, with a minimum of 70 traps per hectare.

When setting up the traps, special attention should be paid to the dynamics of the pest within the property, reinforcing them in places where the greatest pressure is detected.

Hang SWD Lure® 1.5 meters above the ground in the sunny part of the plant; the holes of the trap should not be obstructed by leaves or branches.

SWD Lure® will need to be replenished before it fully evaporates. The time interval for its replacement may vary depending on temperature, humidity, exposure to sunlight, etc.

Crops	Pests	Doses	Application Time
Blueberries, Strawberry, Raspberry	Spot-winged flies (Drosophila suzukii)	A minimum of 70 traps per hectare, depending on the pressure and dynamics of the pest	The time, the concentration and the number of applications will be determined according to the thresholds of economic damage through attractive traps for the detection of adult flies. It is suggested to monitor these traps at least twice a week.
Figs			
Stone fruit trees (cherry tree, peach tree, plum tree, apricot tree)			
Seed fruit trees (vines)			
Tropical fruit trees (mango tree, guava tree, etc.)			

Preservation

Store in a cool and dry place, avoiding exposure to sunlight.

Safety Regulations

Keep out of reach of children and inexperienced people. Handle with gloves. Wash the exposed surface with plenty of soap and water. If ingested, you should induce vomiting and seek medical advice immediately.

Warnings

Avoid product spillages in order to prevent attraction areas outside the trap. Confer with an agricultural science professional before purchasing and using this product.

Susbin S.A. is responsible for the composition of the product until its expiration date.

Susbin S.A. is not responsible for damages of any nature that may derive from a different use of the product.